

PRA-HEP-93/2

THE QUANTUM-DOUBLE FOR A NONSTANDARD DEFORMATION OF A BOREL SUBALGEBRA $sl(2, C)$

Č Burdík^{◊1}, and P. Hellinger^{*2}

**Department of Theoretical Physics and ◊Nuclear Centre,
Faculty of Mathematics and Physics, Charles University,
V Holešovičkách 2, 180 00 Prague 8, Czech Republic*

Abstract

We give a construction of Driinfeld's quantum double for a nonstandard deformation of Borel subalgebra of $sl(2)$. We construct explicitly some simple representations of this quantum algebra and from the universal R-matrix we obtain the explicit solutions of the Yang-Baxter equation in those cases.

PRA-HEP-93/2 hep-th/9303035 March 1993

¹E-mail: Bitnet=(VSETIN AT CSPUNI12)

²E-mail: Bitnet=(HELLINGER AT CSPUNI12)

1 Introduction

One of the most important quantum group constructions is Drienfeld's quantum double [2]. This construction was used by Drienfeld for construction of quantum deformations of simple Lie algebras. In well know example of $U(sl(2))$ we start from a Borel subalgebra and the Drienfeld's quantum double gives $U_q(sl(2)) \otimes U(u(1))$.

It was pointed by Ogievetski [8] that there exists another nonequivalent Hopf algebra deformation of the Borel subalgebra connected via FRT construction [5] with the solution of the Yang-Baxter equation

$$R = \begin{pmatrix} 1 & -\gamma & \gamma & -\gamma\beta \\ 0 & 1 & 0 & \beta \\ 0 & 0 & 1 & -\beta \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (1)$$

which was found in [1] and studied in [3] [4] [6] [7] [10] The aim of this letter is solve an interesting question what we will obtain when the Drienfeld's quantum double construction is applied to this deformation.

2 The basic Hopf algebra and its dual

Let us consider the Hopf algebra $\mathcal{A}=U_\gamma b_-$ over $\mathbf{C}[[\gamma]]$, the formal deformation of $U b_-$ of $sl(2)$ with the generators τ , π and with the relations

$$[\tau, \pi] = -2\pi. \quad (2)$$

The coalgebra structure is given by

$$\begin{aligned} \Delta(\tau) &= \tau \otimes 1 + \Lambda \otimes \tau \\ \Delta(\pi) &= \pi \otimes 1 + \Lambda^{-1} \otimes \pi \\ \epsilon(\tau) &= \epsilon(\pi) = 0 \end{aligned} \quad (3)$$

and antipodes are

$$\begin{aligned} S(\tau) &= -\Lambda^{-1}\tau \\ S(\pi) &= -\Lambda\pi \\ S^\sigma(\tau) &= -\tau\Lambda^{-1} \\ S^\sigma(\pi) &= -\Lambda\pi, \end{aligned} \quad (4)$$

where $\Lambda = (1 - \gamma\pi)^{-1}$.

We may construct the dual Hopf algebra $\mathcal{A}^*$ with the generators T , P and with the pairing

$$\langle T, \tau^n \pi^k \rangle \stackrel{\text{def}}{=} \delta_{n1} \delta_{k0} \quad (5)$$

$$\langle P, \tau^n \pi^k \rangle \stackrel{\text{def}}{=} \delta_{n0} \delta_{k1} \quad (6)$$

From this pairing one may easily deduce the comultiplications

$$\begin{aligned}\Delta(T) &= T \otimes 1 + 1 \otimes T \\ \Delta(P) &= P \otimes e^{2T} + 1 \otimes P\end{aligned}\tag{7}$$

Proof

$$\begin{aligned}\langle \Delta(T), \tau^{n_1} \pi^{k_1} \otimes \tau^{n_2} \pi^{k_2} \rangle &= \langle T, \tau^{n_1} \pi^{k_1} \tau^{n_2} \pi^{k_2} \rangle = \\ &= \delta_{n_1 1} \delta_{k_1 0} \delta_{n_2 0} \delta_{k_2 0} + \delta_{n_1 0} \delta_{k_1 0} \delta_{n_2 1} \delta_{k_2 0}\end{aligned}$$

and

$$\begin{aligned}\langle \Delta(P), \tau^{n_1} \pi^{k_1} \otimes \tau^{n_2} \pi^{k_2} \rangle &= \langle P, \tau^{n_1} \pi^{k_1} \tau^{n_2} \pi^{k_2} \rangle = \\ &= \delta_{n_1 0} \delta_{k_1 0} \delta_{n_2 0} \delta_{k_2 1} + \delta_{n_1 0} \delta_{k_1 1} \delta_{n_2 0} 2^{n_2}\end{aligned}$$

and

$$\langle T^k, \tau^n \rangle = n! \delta_{nk}$$

as will be proven (see the equation (9)). $\square$

We are to find the commutation relations between T and P . The result is

$$[T, P] = -\frac{1}{2} \gamma (e^{2T} - 1)\tag{8}$$

Proof

$$\begin{aligned}\langle TP, \tau^n \pi^k \rangle &= \langle T \otimes P, \Delta(\tau^n \pi^k) \rangle = \\ &= \delta_{n1} \delta_{k1} \\ \langle PT, \tau^n \pi^k \rangle &= \langle P \otimes T, \Delta(\tau^n \pi^k) \rangle = \\ &= \delta_{n1} \delta_{k1} + (n > 0) \gamma \delta_{k0} 2^{n-1},\end{aligned}$$

where the symbol (x) is defined

$$(x) = \begin{cases} 1 & \text{if } x \text{ is true} \\ 0 & \text{otherwise} \end{cases}$$

$\square$

Now is time for counting of powers of the generators. Let us start with promised

$$\langle T^n, \tau^k \pi^m \rangle = n! \delta_{nk} \delta_{m0}\tag{9}$$

Proof

$$\begin{aligned}\langle T^n, \tau^k \pi^m \rangle &= \langle T^{\otimes n}, \Delta^{n-1}(\tau^k \pi^m) \rangle = \\ &= \delta_{m0} \langle T^{\otimes n}, (t_1 + \dots + t_n)^k \rangle\end{aligned}$$

since (5), (2) and $\Lambda = 1 + \gamma\pi + o(\gamma^2)$, where

$$\begin{aligned}t_1 &\stackrel{\text{def}}{=} t \otimes 1 \otimes \dots \otimes 1 \\ t_2 &\stackrel{\text{def}}{=} 1 \otimes t \otimes 1 \otimes \dots \otimes 1 \\ &\vdots \\ t_{n-1} &\stackrel{\text{def}}{=} 1 \otimes \dots \otimes 1 \otimes t \otimes 1 \\ t_n &\stackrel{\text{def}}{=} 1 \otimes \dots \otimes 1 \otimes t\end{aligned}$$

And so $\langle T^{\otimes n}, (t_1 + \dots + t_n)^k \rangle = \delta_{nk} n! \square$

And now we will switch to P

$$\langle P^n, \tau^m \pi^k \rangle = \delta_{m0} (-\gamma)^{n-k} \Theta_k^n \quad (10)$$

where

$$\Theta_k^n = \sum_{p=0}^k \binom{k}{p} (-1)^{k-p} p^n$$

with the following properties

$$\Theta_k^n = 0 \text{ for } k > n \quad \Theta_n^n = n! \quad (11)$$

Proof

$$\begin{aligned}\Delta(\pi) &= \pi \otimes 1 + 1 \otimes \pi - \gamma \pi \otimes \pi \\ \Delta^n(\pi) &= \sum_{i=1}^{n+1} \pi_i + (-\gamma) \sum_{i>j=1}^{n+1} \pi_i \pi_j + (-\gamma)^2 \sum_{i>j>k=1}^{n+1} \pi_i \pi_j \pi_k + \dots + \\ &\quad + (-\gamma)^n \pi_1 \pi_2 \dots \pi_n \pi_{n+1},\end{aligned}$$

by the induction, where π_i is defined similar as for τ . We ought to calculate

$$\begin{aligned}\langle P^n, \tau^k \pi^m \rangle &= \langle P^{\otimes n}, \Delta^{n-1}(\tau^m \pi^k) \rangle = \\ &= \delta_{m0} \langle P^{\otimes n}, \Delta^{n-1}(\pi)^k \rangle\end{aligned}$$

Thanks to (6) we may π_j 's in the previous pairing consider as grassmanian variables $\pi_j^2 = 0$, for $j = 1, \dots, n$ and $\pi_i \pi_j = \pi_j \pi_i$ for $i, j = 1, \dots, n$. Under such ansatz we have this very convenient result

$$\Delta^n(\pi) \stackrel{\langle \cdot \rangle}{=} -\frac{1}{\gamma} (e^{-\gamma \sum_{i=1}^{n+1} \pi_i} - 1) \quad (12)$$

as one may easily check ($\stackrel{\langle \cdot \rangle}{=}$ means the equality in the pairing). So we will proceed

$$\begin{aligned}
\langle P^{\otimes n}, \Delta^{n-1}(\pi)^k \rangle &= \langle P^{\otimes n}, (-\frac{1}{\gamma}(e^{-\gamma \sum_{i=1}^n \pi_i} - 1))^k \rangle = \\
&= \langle P^{\otimes n}, (-\gamma)^{-k} \sum_{p=0}^k \binom{k}{p} (-1)^{k-p} e^{-\gamma p \sum_{i=1}^n \pi_i} \rangle = \\
&= (-\gamma)^{n-k} \sum_{p=0}^k \binom{k}{p} (-1)^{k-p} p^n
\end{aligned}$$

□

Now we may obtain the general formula

$$\langle T^j P^n, \tau^m \pi^k \rangle = m! \delta_{mj} (-\gamma)^{n-k} \sum_{p=0}^k \binom{k}{p} (-1)^{k-p} p^n \quad (13)$$

Proof Straightforward □

At this moment we have the description of the dual Hopf algebra $\mathcal{A}^*$ or precisely we have only the explicit form of the bialgebra structure, but henceforth we will not need the antipodes. Let us construct the double now.

3 The quantum double $D(\mathcal{A})$

We have to combine $\mathcal{A}$ and $\mathcal{A}^\circ$, where $\mathcal{A}^\circ$ is the dual $\mathcal{A}^*$ with the opposite comultiplication. Let us briefly summarize these structures:

$$\begin{aligned}
[\tau, \pi] &= -2\pi \\
[T, P] &= -\frac{1}{2}\gamma(e^{2T} - 1) \\
\Delta(\tau) &= \tau \otimes 1 + \Lambda \otimes \tau \\
\Delta(\pi) &= \pi \otimes 1 + \Lambda^{-1} \otimes \pi \\
\Delta(T) &= T \otimes 1 + 1 \otimes T \\
\Delta(P) &= P \otimes 1 + e^{2T} \otimes P
\end{aligned} \quad (14)$$

For the commutation relations between π , τ and P , T we are forced to use the standard procedure

$$X\zeta = \Sigma(\langle \Delta^2(X), S_1^\sigma \Delta^2(\zeta) \rangle_{13}) \quad (15)$$

for $X \in \{P, T\}$ and $\zeta \in \{\pi, \tau\}$, where $\langle \cdot, \cdot \rangle_{13}$ is the evaluation between the dual algebras on the first and the third position and Σ exchanges $\mathcal{A}^\circ$ and $\mathcal{A}$

This reads in our case as follows:

$$\begin{aligned}
[P, \tau] &= -\gamma\tau - 2P \\
[P, \pi] &= \Lambda^{-1}e^{2T} - 1 + \gamma\pi \\
[T, \tau] &= \Lambda - 1 \\
[T, \pi] &= 0
\end{aligned} \tag{16}$$

Proof

$$\begin{aligned}
P\tau &= \Sigma(\langle e^{2T} \otimes e^{2T} \otimes P + e^{2T} \otimes P \otimes 1 + P \otimes 1 \otimes 1, \\
&\quad S_1^\sigma(\tau \otimes 1 \otimes 1 + \Lambda \otimes \tau \otimes 1 + \Lambda \otimes \Lambda \otimes \tau) \rangle_{13}) = \\
&= \Sigma(\langle e^{2T} \otimes e^{2T} \otimes P + e^{2T} \otimes P \otimes 1 + P \otimes 1 \otimes 1, \\
&\quad -\tau\Lambda^{-1} \otimes 1 \otimes 1 + \Lambda^{-1} \otimes \tau \otimes 1 + \Lambda^{-1} \otimes \Lambda \otimes \tau \rangle_{13}) = \\
&= \Sigma(P\tau - \gamma\tau - 2P) = \\
&= \tau P - \gamma\tau - 2P
\end{aligned}$$

and similiary the others. $\square$

To obtain the classical limit one may introduce the new generators

$$\begin{aligned}
p &\stackrel{\text{def}}{=} \frac{P}{\alpha} \\
t &\stackrel{\text{def}}{=} \frac{T}{\alpha}
\end{aligned} \tag{17}$$

where $\alpha \stackrel{\text{def}}{=} \gamma/2$ In these generators the quantum double $\mathbf{D}(\mathcal{A})$ looks

$$\begin{aligned}
[p, \tau] &= -2\tau - 2p \\
[p, \pi] &= \Lambda^{-1}e^{2\alpha t} - 1 + 2\pi \\
[t, \tau] &= \frac{1}{\alpha}(\Lambda - 1) \\
[t, \pi] &= 0 \\
[\tau, \pi] &= -2\pi \\
[t, p] &= -\frac{1}{\alpha}(e^{2\alpha t} - 1) \\
\Delta(\tau) &= \tau \otimes 1 + \Lambda \otimes \tau \\
\Delta(\pi) &= \pi \otimes 1 + \Lambda^{-1} \otimes \pi \\
\Delta(t) &= t \otimes 1 + 1 \otimes t \\
\Delta(P) &= P \otimes 1 + e^{2\alpha t} \otimes p
\end{aligned} \tag{18}$$

The classical limit is then the enveloping algebra of the Lie algebra L with four generators and Lie brackets have this form:

$$[\tau, \pi] = -2\pi$$

$$\begin{aligned}
[p, \tau] &= -2\tau - 2p \\
[t, p] &= -2t \\
[t, \tau] &= 2\pi \\
[p, \pi] &= 2t \\
[t, \pi] &= 0
\end{aligned} \tag{19}$$

4 The universal R -matrix

We have to find the dual bases for $\mathcal{A}$ and $\mathcal{A}^o$. If define R_j as

$$\begin{aligned}
R_1 &\stackrel{\text{def}}{=} P \\
R_n &\stackrel{\text{def}}{=} P^n - \sum_{k=1}^{n-1} \Theta_k^n P^k (-\gamma)^{n-k}
\end{aligned} \tag{20}$$

we have

$$\langle T^n R_k, \tau^m \pi^l \rangle = n!k! \delta_{nm} \delta_{kl} \tag{21}$$

Proof easy $\square$

Then we have the universal R -matrix in the form

$$\mathcal{R} = \sum_{n,k=0}^{\infty} \frac{1}{n!k!} \tau^n \pi^k \otimes T^n R_k \tag{22}$$

5 Representations of $\mathbf{D}(\mathcal{A})$

The problem to find (all (irreducible)) representations of $\mathbf{D}(\mathcal{A})$ is generally very difficult, but in our case we have luckily at least two representations obtained from the classical limit L :

The two dimensional one ρ , parametrized by the three complex numbers x, y, z

$$\rho_{xyz}(\tau) \stackrel{\text{def}}{=} -\rho_{xyz}(p) \stackrel{\text{def}}{=} \sigma_3 + x\mathbb{1} + y\sigma_- \tag{23}$$

$$\rho_{xyz}(\pi) \stackrel{\text{def}}{=} \rho_{xyz}(t) \stackrel{\text{def}}{=} z\sigma_- \tag{24}$$

where $\mathbb{1}$ is the identity matrix and σ_3, σ_- are Pauli matrices

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad \sigma_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \tag{25}$$

Because $\sigma_-^2 = 0$ the representation ρ of L induce the representation of $\mathbf{D}(\mathcal{A})$ and so we have this solution of YBE:

$$R_{x_1, y_1, z_1, x_2, y_2, z_2} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -\alpha z_1(1+x_2) & 1 & 0 & 0 \\ \alpha z_2(1+x_1) & 0 & 1 & 0 \\ f & \alpha z_2(x_1-1) & -\alpha z_1(x_2-1) & 1 \end{pmatrix} \tag{26}$$

where $f = -\alpha^2 z_1 z_2 (x_1 - 1)(x_2 + 1) + \alpha(z_2 y_1 - z_1 y_2)$. The solution of constant Yang-Baxter equation (setting $x_1 = x_2 = x$, $y_1 = y_2 = y$, $z_1 = z_2 = z$ and $\gamma = \alpha z(x + 1)$, $\beta = \alpha z(x - 1)$)

$$R_{\beta\gamma} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -\gamma & 1 & 0 & 0 \\ \gamma & 0 & 1 & 0 \\ -\gamma\beta & \beta & -\beta & 1 \end{pmatrix} \quad (27)$$

is the transposed (1).

The three dimensional representation we may obtain using the fact that the element $\pi - t$ is central in L and after the factorization $L/(\pi - t)$ we have the 3D Lie algebra $\tilde{L}$ with the structure

$$\begin{aligned} [\tilde{\tau}, \tilde{\pi}] &= -2\tilde{\pi} \\ [\tilde{p}, \tilde{\tau}] &= -2\tilde{\tau} - 2\tilde{p} \\ [\tilde{\pi}, \tilde{p}] &= -2\tilde{\pi} \end{aligned} \quad (28)$$

The representation ϱ is given by the adjoint representation

$$\begin{aligned} \text{ad}\tilde{\pi} &= \begin{pmatrix} 0 & 2 & -2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ \text{ad}\tilde{\tau} &= \begin{pmatrix} -2 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{pmatrix} \\ \text{ad}\tilde{p} &= \begin{pmatrix} 2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & -2 & 0 \end{pmatrix} \end{aligned} \quad (29)$$

$$(30)$$

as follows

$$\begin{aligned} \varrho_{xy}(\pi) &= \varrho_{xy}(t) = y \cdot \text{ad}\tilde{\pi} \\ \varrho_{xy}(\tau) &= \text{ad}\tilde{\tau} + x\mathbb{I} \\ \varrho_{xy}(p) &= \text{ad}\tilde{p} - x\mathbb{I}. \end{aligned} \quad (31)$$

As above since $\text{ad}\tilde{\pi}^2 = 0$ this representation gives the solution of YBE

$$\begin{aligned} R(x_1, y_1, x_2, y_2) &= \mathbb{I} + \alpha(\varrho_{x_1 y_1}(\pi) \otimes \varrho_{x_2 y_2}(p) + \varrho_{x_1 y_1}(\tau) \otimes \varrho_{x_2 y_2}(t)) + \\ &\quad + \alpha^2 \varrho_{x_1 y_1}(\tau\pi) \otimes \varrho_{x_2 y_2}(tp) \end{aligned} \quad (32)$$

6 Conclusion and Remarks

The classical limit of the quantum double $\mathbf{D}(\mathcal{A})$ - Lie algebra L is obviously not $gl(2)$ as one may see after the change of bases in $\tilde{L}$ (28), which provides such

commutation relations:

$$[H, X^\pm] = \pm 2X^\pm \quad [X^+, X^-] = 0 \quad (33)$$

After the finishing of this letter we have obtained the preprint [9], where the author reformulates the FRT method in terms of the Quantum double. He has studied the case of the matrix (1) and he has constructed the corresponding Quantum double (but the formula for the universal R -matrix is given only in several terms of its formal power expansion). It is however generally very difficult to compare those explicit formulae for two algebras. We have solved this problem only for the classical limit and we are able to show the equivalence.

References

- [1] E. E. Demidov, Yu. I. Manin, E. E. Mukhin, D.V.Zhdanovich. Progr. Theor. Phys. Suppl. 102 (1990) 203
- [2] V.G. Drinfeld. *Quantum groups*. In A. Gleason, editor, *Proceedings of the ICM*, pages 798–820, Rhode Island, 1987. AMS.
- [3] H.Ewen, O.Ogievetsky, J.Wess. *Quantum Matrices in Two Dimensions*. Lett. Math. Phys. 22 (1991) 297
- [4] H.Ewen, O.Ogievetsky, J.Wess. *Realizations and Real Forms of Quantum groups in 2 Dimensions*. Preprint MPI-Ph/91-92 München (1991)
- [5] L.D. Faddeev, N.Yu. Reshetikhin, L.A. Takhtajan. *Algebra i Analiz* 1 vol.1 (1989) 178 (in Russian); English translation: Leningrad Math. J. 1 (1990) 193
- [6] J.Hietarinta. Phys.Lett.A 165 (1992) 245
- [7] L.Hlavaty. J.Phys.A – Math.Gen. 25 (1992) L63
- [8] O. Ogievetski. Talk given in the 13th Winter School "Geometry and Physics". Zdíkov, January 9-16, 1993.
- [9] A. A. Vladimirov. JINR preprint E2-93-34 Dubna (1993) (hep-th 9302043)
- [10] S. Zakrzewski. *A Hopf Star-Algebra of Polynomials on the Quantum $SL(2,R)$ a 'Unitary' R -Matrix* Lett. Math. Phys. 22 (1991) 287